

---

IV. Description of a new Species of Lichen. By Dawson Turner,  
Esq. F.R.S. A.S. and L.S.

Read April 17, 1804.

LICHEN PHÆOCEPHALUS.

LICHEN crustâ e granulosâ sub-foliaceâ fuscescente; bacillis nigris; tuberculis rufo-fuscis, margine luteo-albicante.

TAB. VI. Fig. 1.

Innascitur horreorum assibus apud Bruisyard in Suffolciâ, et Lakenham in Norfolciâ.

Crusta latè effusa, crassiuscula, e granulis sub-gelatinosis, satis magnis, incisis, folia simulantibus, madore dilutè viridi-fuscis, siccitate nigricantibus, densè congestis constans; bacilla nigra, lineam ferè alta, crustam vix, aut ne vix superantia, apice in tuberculum sub-rotundum, pulvere rufo-fusco repletum, et margine tenuissimo, albicante, vel luteolo cinctum, tumescientia. Tuberculorum pulveris marginisque color in siccis madidisque idem.

I should not trouble the Linnean Society with a description and figure of this single lichen, were it not for its close affinity to *L. chrysocephalus*, which I described in a former paper\*. I regarded that species as an interesting addition to our Lichenographia; but the present, which I have since been fortunate enough to find in two places, is still more interesting, on account of the singular nature of its crust. In this respect it essentially differs

\* Linn. Trans. vol. vii. p. 88.

from

